

Prevention of VTE Among Hospitalized Patients:

Electronic Alerts
vs. Human Alerts

Background

- Venous thromboembolism (VTE) is a major cause of death, disability and economic burden.
- VTE constitutes the third most common type of cardiovascular illness
- Fatality rate for pulmonary embolism (PE), is approximately 15%. This exceeds the mortality rate for acute myocardial infarction

Background

- Venous thromboembolism (VTE) is often avoidable in hospitalized patients
- Safe and effective prophylaxis for VTE exists, and formal guidelines have been widely disseminated aimed at increasing prophylaxis in high risk patients
- VTE prevention remains underutilized

Background

- At Brigham and Women's Hospital, we have initiated a series of trials aimed at increasing prophylaxis by:
 - **changing MD behavior** and
 - **improving implementation** of prophylaxis

Types of Interventions

- Electronic Computer Generated Mechanism
 - Single Screen Alert
 - Multi-Screen Alert
- Interpersonal Mechanism:
 - Human Alert


First Generation Electronic Alerts

- BWH utilizes BICS (Brigham Integrated Computing System) for all order entry functions
 - Admitting records, demographic information, lab results, medication orders, etc.
- VTE group utilized computer system to screen all patients admitted to the hospital for High Risk VTE Status

First Generation Alert: Development

- Aim: to increase rate of prophylaxis in patients at risk for DVT and PE
- Developed computer program to detect and identify which patients were at risk
- Alert the responsible physician of high risk patient (via e alert) and offer opportunity to order appropriate prophylaxis

Study Schema



DEFINITION OF "HIGH RISK"

VTE risk score \geq 4 points:

- Cancer 3 (ICD codes)
- Prior VTE 3 (ICD codes)
- Hypercoagulability 3 (Leiden, ACLA)
- Major surgery 2 (> 60 minutes)
- Bed rest 1 ("bed rest" order)
- Advanced age 1 (> 70 years)
- Obesity 1 (BMI > 29 kg/m²)
- HRT/OC 1 (order entry)

RANDOMIZATION

**VTE risk score ≥ 4
No prophylaxis
N = 2506**

```
graph TD; A["VTE risk score ≥ 4  
No prophylaxis  
N = 2506"] --> B["INTERVENTION  
Single alert  
n = 1255"]; A --> C["CONTROL  
No alert  
n = 1251"];
```

**INTERVENTION
Single alert
n = 1255**

**CONTROL
No alert
n = 1251**

(Kucher N, et al. NEJM 2005;352:969-977)

Physician Notification of Alerts

The screenshot shows a window titled "Alerts" with a dark red header. Below the header, a grey bar displays "You are None, M.D." with a dropdown arrow. A yellow text box contains the instruction: "There are new alerts on these patients. Mark one and <Enter> to deal with it now, or <Esc> to skip them all." Below this is a list of two alerts:

09:11 AM 10/28	Rxtest, M	DVT HIGH RISK
09:11 AM 10/28	Rxtest, J	DVT HIGH RISK

At the bottom, a grey bar contains the text "<F1> Info. <Esc> Cancel." and two buttons: "OK" and "Cancel".

View PtLookup

Patient: XXXXXXXX,XXXXX 76M 00000000 Adm: 03/02/2005 Room: 8B-312
Time: 03:03 AM Mar 3, 2005 Alert #1881848 8B phone: x7725
Alert: Patient is at high risk for deep vein thrombosis, according to BWH
guidelines.
Reason: Total DVT risk assessment score is 6.
Patient does not have any active Anti-Embolism orders.
Patient is currently NOT on a drug from ANTICOAGULANTS drug family.

Relevant medications and lab results: <alert Details>


Act- [IA Order set: DVT Prophylaxis
ions: [IB Quick Ref: Prevention
[IC Exit to order entry

Covering M.D.: Bp#
<dOne>

<Not my patient> <page M.D.>
<Comments> <Logic >

First Generation Computerized Alerts for VTE prevention

- Utilization of computer generated alerts to house staff reduced the incidence of VTE by 41%
- VTE prophylaxis was prescribed in 33.5% of patients in the intervention group
- Following study conclusion a multidisciplinary team was convened to enhance the VTE alerts at BWH



Kucher N, et al. NEJM. 2005;352:969-977.

CPOE Alerts

- Institutions that utilize decision support and computerized alerts during prescribing have reported high rates of physician override
- A study conducted at BIDMC reported that 94.2% of computerized alerts were overridden
- Reviewers concluded of the 189 rules studied, 36.5% of the rules were invalid and agreed with the physician's decision 97.9% of the time

Saving CPOE from Extinction

- CPOE must evolve to keep up with the growing demand for effective medical informatics and technology solutions
- Next steps are utilizing algorithms that take into account patient specific factors and generate prescribing recommendations to providers

Second Generation: Electronic Computer Generated Alerts

BWH VTE Alerts: The Future

- Goals:
 - Engage the house officer with an interactive alert to increase acceptance and gain feedback
 - Update the DVT prophylaxis template to meet current practice guidelines
 - Provide real-time knowledge links

Interactive Techniques

- Provide objective data to the house officer that this alert positively impacts patient outcome
- Create opportunity to capture rationale for declining alert
 - Hypothesized that many physicians fear a risk of bleeding with anticoagulation
- Provide a final opportunity to order mechanical prophylaxis
- Alert attending physician if alert is not acknowledged after 24 hours

DVT Alert Screen

Time: 03:24 AM Dec 4, 2002 Alert #1014346 14B phone: x7905
Alert: Patient is at high risk for deep vein thrombosis, according to BWH guidelines.

Reason: Total DVT risk assessment score is 4.
Patient does not have any active Anti-Embolism orders.
Patient is currently NOT on a drug from ANTICOAGULANTS drug family.

Relevant medications and lab results: [<alert Details>](#)
Study at BWH published in NEJM 2005;352:969-977 demonstrated a 41% decrease in incidence of VTE using computer generated alerts to House Staff physicians

Act- [IA Order set: DVT PROPHYLAXIS TEMPLATE.
ions:[IB Partners Handbook: VTE Guidebook 4th edition
[IC Exit to order entry

Covering M.D.: Bp#
<dOne>

<Not my patient>

<pAge M.D.>

<coMments>

< Logic >



QA

Rule Logic – Alert Details

Time: 03:22 AM Nov 2, 2004 Alert #1740559 7A phone: x7695

Details for alert #1740559

Rule: Patient is at high risk for deep vein thrombosis, according to BWH guidelines.

The following risk factors were found to be positive (score applied):

Patient is overweight: BMI >29 kg/m² (1)

has history of deep vein thrombosis or pulmonary embolism (3).

-and-

Patient does not have any active Anti-Embolism orders.

-and-

Patient is currently NOT on a drug from ANTICOAGULANTS drug family.

Option A

Time: 03:24 AM Dec 4, 2002 Alert #1014346 14B phone: x7905
Alert: Patient is at high risk for deep vein thrombosis, according to BWH guidelines.

Reason: Total DVT risk assessment score is 4.
Patient does not have any active Anti-Embolism orders.
Patient is currently NOT on a drug from ANTICOAGULANTS drug family.

Relevant medications and lab results: [<alert Details>](#)

Study at BWH published in NEJM 2005;352:969-977 demonstrated a 41% decrease in incidence of VTE using computer generated alerts to House Staff physicians

Act- [1A Order set: DVT PROPHYLAXIS TEMPLATE.
ions:[1B Partners Handbook: VTE Guidebook 4th edition
[1C Exit to order entry



Covering M.D.: Bp#
[<dOne>](#)

[<Not my patient>](#)

[<pAge M.D.>](#)
[<coMments>](#)

[< Logic >](#)



QA

ViewOrders PtLookup Feedback Help Goodbye

OETEST,CLOVIS 32F 11489945

Adm: 11/01/91 Room: 17A-444

DVT Prophylaxis Order Set Page 1

Consider combined pharmacological and mechanical prophylaxis in high-risk patients.

MECHANICAL PROPHYLAXIS

Select one or more.

- A Anti-embolism - TED stockings KNEE HIGH
- B Anti-embolism - TED stockings THIGH HIGH
- C Anti-Embolism - Pneumatic Compression - CALF ONLY
- D Anti-Embolism - Pneumatic Compression - CALF and THIGH

PHARMACOLOGICAL PROPHYLAXIS

Select only one.

- E ENOXAPARIN 40 MG SC QD
- F ENOXAPARIN 30 MG SC QD for patients with renal impairment
- G HEPARIN 5,000 UNITS SC TID
- H HEPARIN 5,000 UNITS SC BID
- I FONDAPARINUX 2.5 MG SC QD
- J FONDAPARINUX 2.5 MG SC QOD For patients with renal impairment

OK

Cancel

Option B

Time: 03:24 AM Dec 4, 2002 Alert #1014346 14B phone: x7905


Alert: Patient is at high risk for deep vein thrombosis, according to BWH guidelines.

Reason: Total DVT risk assessment score is 4.
Patient does not have any active Anti-Embolism orders.
Patient is currently NOT on a drug from ANTICOAGULANTS drug family.

Relevant medications and lab results: [<alert Details>](#)

Study at BWH published in NEJM 2005;352:969-977 demonstrated a 41% decrease in incidence of VTE using computer generated alerts to House Staff physicians

Act- [JA Order set: DVT PROPHYLAXIS TEMPLATE.
ions:[JB Partners Handbook: VTE Guidebook 4th edition
[JC Exit to order entry



Covering M.D.: Bp#

[<dOne>](#)

[<Not my patient>](#)

[<pAge M.D.>](#)

[<coMments>](#)

[< Logic >](#)



TOA

 Print this page
  Add to my links
  My links
  Modify my links
  Help

Venous Thromboembolism Guidebook 4th Edition

Introduction

Brigham and Women's Hospital.

Contacts:

Eli V. Gelfand, MD,
Gregory Piazza, MD,
Samuel Z. Goldhaber, MD.

Last updated: June 2003


For further information contact Samuel Z. Goldhaber, MD at:
sgoldhaber@partners.org

Introduction

This Venous Thromboembolism Guidebook incorporates evolving contemporary concepts in diagnosis and management of pulmonary embolism (PE) and deep venous thrombosis (DVT) into a user-friendly menu. The purpose of this document is to provide a literature-based review of the current clinical approach to venous thromboembolism as well as up-to-date references for further study in this important topic.

PE Support Group

Table of Contents

	Introduction
	Risk Factors for Thromboembolism
	Diagnosis
	Diagnosis of PE
	Diagnosis of DVT
	Risk Stratification of PE
	Management of PE and DVT
	Management of PE
	Management of DVT
	Therapy of DVT and PE
	Anticoagulation
	IVC Filters
	Thrombolysis
	Embolectomy
	Prevention
	BWH Venous Thromboembolism Guidebook - PDF

Option C or "Done"

Time: 03:24 AM Dec 4, 2002 Alert #1014346 14B phone: x7905

Alert: Patient is at high risk for deep vein thrombosis, according to BWH guidelines.

Reason: Total DVT risk assessment score is 4.
Patient does not have any active Anti-Embolism orders.
Patient is currently NOT on a drug from ANTICOAGULANTS drug family.

Relevant medications and lab results: [Alert Details](#)

Study at BWH published in NEJM 2005;352:969-977 demonstrated a 41% decrease in incidence of VTE using computer generated alerts to House Staff physicians

Act- [JA Order set: DVT PROPHYLAXIS TEMPLATE.
ions:[JB Partners Handbook: VTE Guidebook 4th edition
[JC Exit to order entry

Covering M.D.: Bp#

<Done>

<Not my patient>

<page M.D.>

<Comments>

< Logic >



QA

View PtLookup

Reason for Declining VTE Prophylaxis

Study at BWH published in NEJM 2005;352:969-977
demonstrated a 41% decrease in incidence of VTE using
computer generated alerts to the House Staff physicians

Provide Reason for Declining VTE Prophylaxis

- Patient already receiving anticoagulants
- Risk of bleed outweighs benefit of anticoagulant therapy
- Patient is "Comfort Measures Only"
- Scheduled procedure
- Other

Other reason (minimum of 15 characters)

|

|

Done



Mechanical DVT Prophylaxis Page 1

There is not an increased risk of bleed with mechanical prophylaxis such as graduated compression stockings or intermittent pneumatic compression devices. Consider ordering one or more of the following mechanical prophylaxis orders.

Select one or more:

- A Anti-embolism - TED stockings KNEE HIGH
- B Anti-embolism - TED stockings THIGH HIGH
- C Anti-embolism - Pneumatic boots CALF ONLY
- D Anti-embolism - Pneumatic boots CALF AND THIGH

Edit

OK

Cancel

DEV

Escalation and Timing of Alerts

- Alerts should be set up to generate each day at 8:30 AM
- If an alert was not acknowledged after 24 hours the attending physician on record should be text paged.

Date	Time	Recipient	Message
10/28/2005	02:37:11PM	Goldhaber, Samuel Zachary, MD	8888-19327139 MRN: 19327139 LOC: 17A-301 High risk VTE patient without prophylaxis orders.
10/28/2005	02:36:08PM	Goldhaber, Samuel Zachary, MD	8888-19327147 MRN: 19327147 LOC: 17A-311 High risk VTE patient without prophylaxis orders.

Quality Assurance

- Weekly reports are reviewed
- Allows core team to review all aspects of the alerts including:
 - type of action taken
 - rate of overrides
 - reasons for declining the alerts

VTE Prophylaxis: eALERT Perspectives

- eAlert Trial: lead article in March 10, 2005 NEJM, with accompanying Editorial and CME Exam, receives widespread attention.
- Many hospitals don't have comparable IT to replicate BWH.
- Question arises: Are BWH results applicable to other USA hospitals ?

Human Alerts

VTE Prophylaxis: hALERT

- We have initiated a multicentered RCT of human alerts (hALERT).
- **Objective:** to recruit hospitals that differ from BWH re: IT, community vs. academic, urban vs. suburban/rural, location within USA.
- Can a **human** alert be more effective than an **electronic** alert?

Methodology

- Patients admitted to the hospital are screen by human for increased VTE risk
- High risk patients are randomized to alert or no alert
- Physicians of patients in alert group receive page alerting them of High risk status
- Records are checked for prophylaxis order 48 hours after alert
- 90 day follow up for clinically significant VTE and clinically importing bleeding

hALERT Screeners: Require Backup

- The study must be done on weekdays, non-Holidays, at a minimum.
- There must be backup screeners, in case the principal screener is ill or is on vacation.

hALERT: Capturing new prophylaxis orders

- Enrolled patients must be reexamined in 24-48 hours to determine whether prophylaxis orders were written.
- Capturing prophylaxis orders after enrollment applies to both the Intervention Group and to the Control Group.

HUMAN ALERT TRIAL

1. Human (often RN or pharmacist) issues the Alert, not a computer
2. The Attending Physician, not the Intern, receives the Alert
3. Diversity of centers: community, suburban, throughout the USA
4. Will Attendings pay more attention than House Staff?

HUMAN ALERT TRIAL

- 900 patients were enrolled from 10 institutions within the first 28 weeks of the trial (as of January 5, 2007)
- Geographic diversity: from Massachusetts to Salt Lake City
- Hospital diversity: community, academic, VAMC
- Weekly eNewsletter
- Recruiting additional sites

Conclusions

- Changing physician behavior is challenging
- Multi-disciplinary team involvement is critical to successful implementation
- Need to engage providers and obtain feedback
- Designing “smart alerts” that include decision support functionality or “human alerts” that require face to face contact may be effective

Thank you for your attention!
